


Electrolytic Cells


- Voltaic cells are spontaneous redox reactions which convert chemical energy to electrical energy. Voltaic cells are essentially batteries. *requires a battery*
- * • A cell that uses an external source of electrical energy to drive a non-spontaneous redox reaction is called an electrolytic cell
- Electrolysis is the process that takes place in a electrolytic cell and it literally means to "break apart"
- * • Since electrolytic cells are non-spontaneous, the reducing agent must be higher than the oxidizing agent on the table on pg. 7 of data booklet (*weak agents*)
- Let's compare a voltaic cell to an electrolytic cell



voltaic cell

- Zn is SRA
- Zn has more "-" electric potential (E°)

- Cu^{2+} is SOA
- Cu^{2+} has more "+" electric potential (E°)



electrolytic cell

- Zn^{2+} is the more "-" E°
- Zn^{2+} is weak OA

- Cu(s) is the more "+" E°
- Cu(s) is weak RA

- The cell potential (E°_{cell}) for an electrolytic cell is calculated the same way as a voltaic cell. All voltaic cells will have a positive cell potential and all electrolytic cells will have a negative cell potential.

voltaic $E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}}$

$$E^\circ_{\text{cell}} = 0.34\text{V} - (-0.76\text{V})$$

$$E^\circ_{\text{cell}} = 1.10\text{V}$$

Electrolytic

$$E^\circ_{\text{cell}} = (-0.76\text{V}) - (0.34\text{V})$$

$$E^\circ_{\text{cell}} = -1.10\text{V}$$

- Comparing a voltaic cell to an electrolytic cell

Voltaic Cell	Electrolytic Cell
spontaneous	non-spontaneous
converts chemical energy to electrical energy	converts electrical energy to chemical energy
is a battery	requires a battery
anions move to anode and cations move to cathode	anions move to anode and cations move to cathode
electrons flow into cathode	electrons flow into cathode
<ul style="list-style-type: none"> ▪ oxidation at anode ▪ more negative electrical potential (E°) at anode ▪ strongest reducing agent at anode 	<ul style="list-style-type: none"> ▪ oxidation at anode ▪ more positive electrical potential (E°) at anode ▪ weakest reducing agent at anode
<ul style="list-style-type: none"> ▪ reduction at cathode ▪ more positive electrical potential (E°) at cathode ▪ strongest oxidizing agent at cathode 	<ul style="list-style-type: none"> ▪ reduction at cathode ▪ more negative electrical potential (E°) at cathode ▪ weakest oxidizing agent at cathode
positive cell potential (E°_{cell})	negative cell potential (E°_{cell})

*****Now try Practice Problems*****


Practice Problems: Electrolytic Cells

1. Consider the following un-balanced redox reactions:
 - a. $I_2 + Cl^- \rightarrow I^- + Cl_2$
 - b. $MnO_4^-(aq) + Br^-(aq) \rightarrow Br_2(l) + Mn^{2+}(aq)$ (acidic conditions)



For each un-balanced redox reaction above:

- i. indicate if the reaction is spontaneous or not
 - ii. write out the half-reaction at the anode
 - iii. write out the half-reaction at the cathode
 - iv. write out the balanced redox reaction
 - v. calculate the cell potential
 - vi. indicate if it is a redox reaction that can take place in a voltaic cell or electrolytic cell
2. Oxidation takes place at what electrode in a voltaic cell?
 3. Oxidation takes place at what electrode in an electrolytic cell?
 4. An electrolytic cell and a voltaic cell can look very similar except for the presence or absence of what?
 5. In a voltaic cell, electrons flow out of which electrode?
 6. In an electrolytic cell, electrons flow out of which electrode?
 7. Why does a voltaic cell have the strongest oxidizing agent at the cathode while an electrolytic cell has the weakest oxidizing agent at the cathode?
 8. Consider an electrolytic nickel-cadmium cell.
 - i. Identify the anode and cathode.
 - ii. Write out the oxidation half-reaction, the reduction half-reaction, and the net redox reaction.
 - iii. Calculate the cell potential.
 9. Consider a voltaic nickel-cadmium cell.
 - i. Identify the anode and cathode.
 - ii. Write out the oxidation half-reaction, the reduction half-reaction, and the net redox reaction.
 - iii. Calculate the cell potential.

Electrolytic Cells
Practice Problems
(Solutions)



I.) non-spontaneous






v.) $E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}}$

$E^\circ_{\text{cell}} = +0.54\text{V} - (+1.36\text{V}) = \boxed{-0.82\text{V}}$

VI.) Electrolytic cell

b.) I.) spontaneous



v.) $E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}}$

$E^\circ_{\text{cell}} = +1.51\text{V} - (-1.07\text{V}) = \boxed{0.44\text{V}}$

VI.) Voltaic cell

2.) anode

3.) anode

4.) an electrolytic cell requires a battery/power source

5.) anode

6.) anode

7.) In both a voltaic and an electrolytic cell, the oxidizing agent needs to be at the cathode (i.e. where reduction occurs). An electrolytic cell requires a weak OA so that no spontaneous redox rxns occur. A voltaic cell requires a strong OA so that a spontaneous rxn occurs.

8.) i.) anode is Ni(s) ; cathode is Cd(s)

ii.) oxidation: $\text{Ni(s)} \rightarrow \text{Ni}^{2+}(\text{aq}) + 2\text{e}^-$

reduction: $\text{Cd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cd(s)}$

net: $\text{Ni(s)} + \text{Cd}^{2+}(\text{aq}) \rightarrow \text{Ni}^{2+}(\text{aq}) + \text{Cd(s)}$

iii.) $E^\circ_{\text{cell}} = -0.40\text{V} - (-0.26\text{V})$

$$E^\circ_{\text{cell}} = -0.14\text{V}$$

9.) i.) anode is Cd(s) ; cathode is Ni(s)

ii.) oxidation: $\text{Cd(s)} \rightarrow \text{Cd}^{2+}(\text{aq}) + 2\text{e}^-$


reduction: $\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni(s)}$

net: $\text{Cd(s)} + \text{Ni}^{2+}(\text{aq}) \rightarrow \text{Cd}^{2+}(\text{aq}) + \text{Ni(s)}$




iii.) $E^\circ_{\text{cell}} = -0.26\text{V} - (-0.40\text{V})$

$$E^\circ_{\text{cell}} = +0.14\text{V}$$

- If electrolysis means to break apart, let's consider the electrolysis of water. The equation that represents the breaking apart of water is as follows:



The half-reactions that are involved are found on the table in pg. 7 of the data book based off the reactants and products of the electrolysis process.





Calculation for cell potential:

$$E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}} = -0.83\text{V} - (+1.23\text{V})$$

$$E^\circ_{\text{cell}} = -2.06\text{V}$$

* not spontaneous




↳ twice as much as oxygen!



→ liquid form

EXAMPLE: Consider the electrolysis of molten sodium chloride.

The equation that represents the breaking apart of $\text{NaCl}_{(l)}$ is as follows:



The half-reactions that are involved are:




* not spontaneous due to spontaneity rule!

Calculation for cell potential



$$E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}}$$

$$E^\circ_{\text{cell}} = -2.71 \text{ V} - (+1.36 \text{ V})$$

$$E^\circ_{\text{cell}} = -4.07 \text{ V}$$




- We need to be careful with the electrolysis of aqueous solutions. Consider the electrolysis of aqueous sodium chloride, $\text{NaCl}_{(aq)}$.
 - If we know electrolysis is the process of "breaking down" we would expect the half-reactions for this process to be as follows:




- * ◦ This would be incorrect because we are ignoring the presence of water and the fact that water is both an oxidizing and reducing agent

- We first need to list all substances present and identify them as oxidizing or reducing agents. We will use the agents in half-reactions to create the net redox reaction. The main redox reaction is the reaction that requires the least cell potential/voltage.



Possibility #1



$\text{H}_{2(\text{g})}$ & $\text{Cl}_{2(\text{g})}$
get produced!






$$E^\circ_{\text{cell}} = -0.83\text{V} - (+1.36\text{V})$$

$$E^\circ_{\text{cell}} = -2.19\text{V}$$

* main rxn as it requires the least cell potential!

Possibility #2



~~$$E^\circ_{\text{cell}} = -2.71\text{V} - (+1.23\text{V})$$~~

~~$$E^\circ_{\text{cell}} = -3.94\text{V}$$~~

Helpful Hint to Determine the Anode and Cathode

VOLTAIC CELL

- **Cathode:** reduction half-reaction/ strong oxidizing agent → therefore, needs a half-reaction with a more positive reduction potential (E°)
- **Anode:** oxidation half-reaction/ strong reducing agent → therefore, needs a half-reaction with a more negative reduction potential (E°)

ELECTROLYTIC CELL (just the opposite rules)

- **Cathode:** reduction half-reaction/ weak oxidizing agent → therefore, needs a half-reaction with a more negative reduction potential (E°)
- **Anode:** oxidation half-reaction/ weak reducing agent → therefore, needs a half-reaction with a more positive reduction potential (E°)